

Table 8. Estimated numbers of persons living with HIV/AIDS, by year and selected characteristics, 2001–2004—35 areas with confidential name-based HIV infection reporting

	2001	2002	2003	2004
Age as of end of year (yrs)				
<13	4,838	4,598	4,187	3,713
13–14	830	950	1,123	1,239
15–19	2,745	2,968	3,301	3,683
20–24	11,500	12,033	12,582	13,371
25–29	25,334	25,768	26,538	27,836
30–34	52,672	51,896	50,416	49,133
35–39	82,539	83,133	81,891	80,086
40–44	79,325	86,422	93,414	99,742
45–49	59,008	65,879	72,498	79,728
50–54	34,929	40,488	45,932	52,658
55–59	16,271	19,550	23,286	27,571
60–64	7,781	9,497	11,098	13,170
≥65	6,674	7,816	9,099	10,861
Race/ethnicity				
White, not Hispanic	133,475	141,120	148,459	157,172
Black, not Hispanic	181,964	195,147	206,936	220,028
Hispanic	63,105	68,252	72,967	78,039
Asian/Pacific Islander	1,796	2,084	2,415	2,765
American Indian/Alaska Native	1,587	1,728	1,865	1,996
Transmission category				
Male adult or adolescent				
Male-to-male sexual contact	159,937	172,502	184,778	199,085
Injection drug use	57,287	58,959	60,113	61,799
Male-to-male sexual contact and injection drug use	21,643	22,201	22,688	23,337
Heterosexual contact	34,386	37,986	41,291	44,655
Other ^a	3,445	3,524	3,602	3,702
Subtotal	276,698	295,172	312,472	332,578
Female adult or adolescent				
Injection drug use	30,977	32,003	32,742	33,621
Heterosexual contact	68,173	74,925	81,007	87,262
Other ^a	2,077	2,236	2,383	2,523
Subtotal	101,227	109,164	116,133	123,405
Child (<13 yrs at diagnosis)				
Perinatal	5,810	5,954	6,057	6,100
Other ^b	706	703	698	704
Subtotal	6,515	6,657	6,755	6,804
Total^c	384,446	410,998	435,364	462,792

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 2000, the following 35 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, Guam and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^b Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^c Includes persons of unknown race or multiple races and persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.